

ORIGINAL RESEARCH ARTICLE

Synthesis and biological studies of some novel furo naphthyridine compounds contain chalcone

Fatimah Nazar Mahmood¹, Alaa I. Ayoo², Shakir M. Saied³, Mohanad Y. Saleh^{4,*}

¹ Technical Research Center, Northern Technical University, Mosul, 41001, Iraq

² Department of New and Renewable Energies, College of Science, University of Mosul, 42001, Iraq

³ College of Pharmacy, Al-Noor University, Mosul city, Alshallalat Road, Mosul, 41001, Iraq

⁴ Department of Chemistry, College of Education for Pure Science, University of Mosul, Mosul, 42001, Iraq

*Corresponding author: Mohanad Y. Saleh, mohanadalallaf@uomosul.edu.iq

ABSTRACT

Through a Vilsmeier–Haack condensation 2-chloro-3-formyl-1,8-naphthyridine (I) was synthesized. When treated with aqueous hydrochloric acid converted to compound (II) was successfully obtained in high yield and then converted to Novel 1-(8-methyl) furo[2,3-b]-(1,8-naphthyridine-2-yl) ethenone (III) through Claisen–Schmidt condensation. The condensation between compound III (ketone) and benzaldehyde yield Novel (furo[2,3-b] (1,8-naphthyridine-2-yl)-2-acetyl called chalcone (4). The reaction between chalcone and bromine water yields dibromide (6), Iodo chalcone (5) produced by treatment of chalcone with one to two pieces of crystal iodine in dimethyl sulfoxide. Spectral analysis techniques such as ¹H-NMR, FT-IR were employed to identify and confirm the structural formula of all products. The synthesized compounds (3-6) were evaluation for their anti- bacterial activity against both gram negative and gram-positive bacteria, demonstrating significant compared to standard drug. All these compounds showed moderate activity.

Keywords: furo2,3-b; chalcone derivatives; Vilsmeier–Haack reaction; Claisen–Schmidt condensation; antibacterial activity; dibromide; iodo chalcone

ARTICLE INFO

Received: 17 November 2025

Accepted: 19 December 2025

Available online: 26 March 2026

COPYRIGHT

Copyright © 2026 by author(s).

Applied Chemical Engineering is published by Arts and Science Press Pte. Ltd. This work is licensed under the Creative Commons

Attribution-NonCommercial 4.0 International License (CC BY 4.0).

<https://creativecommons.org/licenses/by/4.0/>

Highlight

- Novel furo2,3-b chalcone derivatives were synthesized via multi-step reactions.
- Structures were confirmed by IR and ¹H NMR spectroscopy.
- Halogenated derivatives (dibromo and iodo chalcones) were successfully prepared.
- Compounds 3–6 exhibited moderate antibacterial activity against Gram-positive and Gram-negative bacteria.
- The results suggest these scaffolds as promising leads for antimicrobial drug development.

1. Introduction

Naphthyridine compounds have played a significant role in heterocyclic chemistry, where their mixed acid-base properties have altered the expected products as they react with electrophilic and cyclophilic agents, and through the carbon and nitrogen atoms. The chemistry of naphthyridine compounds have been of increasing interest since these compounds have been found to be useful as

chemotherapeutic agent against malaria^[1], bacteria^[2] and widely distributed in nature as essential to life fund as potential^[3]. The hetero cyclic compounds that contain nitrogen and oxygen are one of the most extensive compounds that show diverse pharmacological activities. The chalcones derivatives of these groups show activities as anti-tuberculosis agent^[4,5] as important as natural products belonging to flavonoid family^[6]. Chalcones are precursors in the preparation of many important hetero cyclic such as flavones^[7], benzo thiazepine^[8] pyrazolines^[9]. Diels-Alder cyclization was the most important method used to synthesis naphthyridine compounds^[10] 1,8-naphthyridine and tri ethylene glycol ether-Linked are used as important binding unit in molecular design of synthetic receptors^[11] and the compound fluoro benzyl)-3-(2-tolyl)-1,8-naphthyridine (1H)-2-one are used in memory treatment in Al-Zheimer's disease^[12]. Now a day, the quinoline compounds are used as starting materials for synthesis of many fused naphthyridine compounds^[13,14] new dibromide and iodo naphthyridine compounds have been prepared through selective iodination and bromination of chalcones^[15]. Bacteria and fungi that are resistant to antibiotics require ongoing research and development of new antibacterial agents through the synthesis of novel chemicals^[16]. It also possesses many therapeutic properties and a wide spectrum of antibacterial, antifungal and antiviral properties, and has played an important role in many cancer treatments and chronic diseases. In our work, a mouse ring was synthesized on synthetic naphthyridine via the Vilsmeier–Haack reaction, which contained a terminal acetyl group at position 2 of the furan ring, which was converted to a chalcone, and the products had good biological effects. This study aimed to synthesize novel furonaphthyridine derivatives and characterized and identify by using IR, ¹H-NMR and then studies their antibacterial activity against Gram positive and negative bacteria.


2. Experimental

The melting point was measured on electro-thermal CA9300 apparatus and uncorrected. The infrared were measured on IR spectrophotometer model Tensor 27, Bruker Co., Germany by using KBr aiscs. The ¹HNMR are used identify the structure on a JEol 400MHZ, using TMS as internal standard.

Synthesis of 2-chloro-3-formyl -1,8-naphthyridine (1)^[17,18]

A solution of N-(pyridine-2-yl) acetamide


A round-bottom flask with a dropping funnel filled with (14 mol) of POCl₃ filled with a mixture of (2 mol) and 6 mol of dry dimethyl form amide. The circular was submerged in an ice bath between 0 and 50 degrees Celsius. After maintaining the temperature, POCl₃ was added drop by drop while stirring continuously. The dropping funnel was then removed, and the reaction mixture was refluxed at 80 to 900 degrees Celsius for 16 hours. The mixture was then poured into cold water and stirred for one hour, after which the precipitation was filtered, cleaned with cold water, and the crystals were recrystallized using methanol.



The yield was 60%, the m.p. was determined to be between 164 and 166°C, and the hue was light yellow. IR (KBr, cm⁻¹): 3055 (Ar-H), 2720 (CHO), 1685 (C=O), 1565 (C=N), 775 (C-Cl), ¹HNMR (400 MHz, DMSO-d₆, ppm) 10.54 (s, 1H, CHO), 8.45 (d, 1H, C7), 7.86 (s, 1H, C4), 7.45 (d, 1H, C5), and 7.27 (t, 1H, C6).

Synthesis of 2-hydroxy-3-formyl-1,8-naphthyridine (2)


A mixture of compound (1) (0.01mol) and (35mL, 4mol) of aqueous hydrochloric acid was heated and reflux for (6 hrs.) then allowed to cool to room temperature. The mixture was poured onto ice cold water; the solid was separated and recrystallized from aqueous acetic acid.



The yield was 70% yellow, and the m.p. was determined to be between 151 and 153°C. IR (KBr, V, cm^{-1}): 1690(C=O), 1665(C=N), 3065 (Ar-H) and 3230(OH). ^1H NMR (400MHz, DMSO- d_6 , δ , ppm): 8.77(s, 1H, C=O), 8.21(d, 1H, C7), 8.05(s, 1H, C4), 7.18(d, 1H, C5), 7.02(m, 1H, C6) and 15.6(s, 1H, OH).

Synthesis of furo[2,3-b](1,8-naphthyridine-2-yl)-2-acetyl (3)


Compound (2) (0.1mol) chloroacetone (0.1mol) and anhydrous potassium carbonate (0.1mol) in (80ml) of dimethyl formamide was stirred for (6hrs.) at room temperature. Then mixture was reflux on water bath for 24hrs. to (70-90 °C) Then allowed to cool at room temperature then the mixture was poured into crush ice, then precipitate was filtered dried and recrystallized by used DMF.



The m.p was found (125-127°C), the yield 50% yellow brown. IR(KBr, V, cm^{-1}): 1710(C=O), 2976(CH), 1655(C=N) and 1120(C-O-C), ^1H NMR (400MHz, DMSO- d_6 , δ , ppm). 2.99(s, 3H, CH₃), 6.99(s, 1H, furane), 7.15(t, 1H, C6), 8.07(s, 1H, C4), 8.21(m, 1H, C5) and 8.78(d, 1H, C7).

Synthesis of 1-furo[2,3-b](1,8-naphthyridine-2-yl)ethanone derivative (chalcone) (4)^[19]


A mixture of compound (3)(0.01mol) and benzaldehyde (0.01mol) in ethanol (70mL) in the presence of aqueous potassium hydroxide (40%, 15mL) is stirred at room temperature for (24hrs.). Then the reaction mixture was cooled for (10hrs.) and then poured into (300mL) ice cold water, acidified with dilute HCl, the solid was separated then air dried and recrystallized using ethanol.



The m.p found(178-180°C), the yield (70%) color pale brown. IR(BKBr, V, cm^{-1}): 1685(C=O), 1650(C=N), 1565(C=C), 1120(C-O-C) and 1650(C=N). ^1H NMR (400MHz- d_6 , δ , ppm) 6.53(d, CH, ethylene), 7.23-7.40(m, 6H, CH furane benzene ring), 8.11(s, 1H, C4), 8.67-8.68(d, 1H, C5), 7.30-7.32(d, 1H, C7) and 7.41-7.43(t, 1H, C6).

Synthesis of dibromide (5)^[20,21]


Bromine solution (0.1mol) was added dropwise over 30 minutes to cold solution of compound (4) in (15mL) of dichloromethane the mixture stirred overnight, then the solvent reduced to half by evaporator. The precipice was collected and recrystallized by using chloroform.



The m.p. is (203-205°C), color brown, the yield 70%, IR (KBr, V, cm^{-1}), 1695(C=O), 1650(C=N), 1125(C-O-C), 3225(Ar-H). ^1H NMR(400MHz, DMSO- d_6 , δ , ppm). 4.20-4.32(m, 2H, 2-furane, CH-Br), 6.71(d, 1H, CH), 7.59-7.92(m, 6H, benzene, C6), 7.92(d, 1H, C5), 8.56(s, 1H, C4), 7.48-7.49(d, 1H, C7).

General procedure for synthesis of iodo chalcones(6)^[22,23]

Two drops of sulfuric acid were added to the combination of chalcone (4) (3 mmoles) in 15 mL of DMSO after an iodine crystal (1,2) was added. After the mixture refluxed for five hours, it was dumped into cold water and agitated for two hours. The precipitate was filtered, cleaned with a five percent sodium thiosulphate solution, and then recrystallized using ethanol and twenty-five milliliters of cold water.



The m.p (227-229⁰C), yield (50%), color yellow. IR (KBr, V, cm⁻¹): 1668(C=O), 1495(C=C), 1560(C=N) and 3150(Ar-H). ¹HNMR (400MHZ, DMSO-d₆-δ, ppm). 7.23-7.55(m,8H, phenyl ring, 2-furan, CH=, C6), 8.15(s,1H, C4), 8.23(d, 1H, C5) and 8.52(d, 1H, C7).

All synthesized compounds are shown in **Figure (1)** & The physical and spectral characteristics are shown in **Tables (1-2)**.




Figure 1. Synthesis of furonaphthyridine derivatives (I-VI).

3. Result and discussion

Naphthyridine may be made by using a variety of techniques, although the Vilsmeier approach is the most crucial for creating substituted naphthyridine^[24,25]. In this study, 2-chloro-3-formyl-1,8-naphthyridine (1) was made by reacting to N-(pyridine-2-y)-acetamide with Vilsmeier reagent. POCl₃ was added to the substrate in DMF at a temperature between 0 and 50 degrees Celsius and refluxing it for one hour. The **Figure (2)** describes how the response mechanism operates^[26]:




Figure 2. The mechanism of formation vilsmeier – Haack reagent.

Compound (1) IR spectra revealed a distinctive absorption band at 1685 cm^{-1} for $\text{C}=\text{O}$, 1565 cm^{-1} for $\text{C}=\text{N}$, and 775 cm^{-1} for $\text{C}-\text{Cl}$. The molecule's (1) ^1H NMR spectra revealed the following notable peaks: a singlet band at 10.54 ppm for aldehyde proton (CHO), at 7.86 ppm for H-4, a doublet at 8.45 ppm for H-7, at 7.45 ppm for H-5, and a triplet at 7.27 ppm for H-6 table(1,2). Compound (2) was created by refluxing these chemicals (1) in diluted aqueous hydrochloric acid. The IR spectrum for these compounds showed a characteristic band at 1690 cm^{-1} for carbonyl group. The ^1H NMR for compound (2) showed a singlet at 8.77 ppm for (CHO) group and at 5.60 ppm for OH group and between ($7.02\text{--}8.70$) for proton of naphthyridine ring table(1,2). The compound (3) was prepared from reaction of compound (2) with chloro acetone and potassium carbonate. The IR spectra for these compound showed many band at 1710 cm^{-1} for ($\text{C}=\text{O}$) group and at 1655 cm^{-1} ($\text{C}=\text{N}$) group and at 1120 cm^{-1} for ($-\text{C}-\text{O}-\text{C}-$) group. The ^1H NMR spectra for these compounds showed three singlet signal at 2.99 ppm for CH_3 group and at 6.99 ppm for furane proton and at 8.07 ppm for C4 proton for naphthyridine ring table(1,2)^[27]. Clasen-Schmidt condensation^[28] used to prepared compound (4) from reaction of compound (3) with benzaldehyde in ethanolic solution^[19,20]. These compounds' (4) infrared spectra revealed a prominent band at 1685 cm^{-1} for the ($\text{C}=\text{O}$) group and 1565 cm^{-1} for the ($\text{C}=\text{C}$) group. The ^1H NMR spectra of compound (4) revealed three signals: a doublet at 6.53 ppm , $8.67\text{--}8.68\text{ ppm}$, $7.30\text{--}7.32\text{ ppm}$ for the $\alpha\text{-CH}$ proton of ethylene and for ($\text{C}5,\text{C}7$) for the naphthyridine ring, a singlet at 8.11 ppm for the C4 proton for the naphthyridine ring, a multiplet at $7.23\text{--}7.40\text{ ppm}$ for the benzene proton and CH Furane, and a triplet at $7.41\text{--}7.43$ for the naphthyridine ring table(1,2). Compound (4) was brominated to produce dibromide (5).

The reaction done by adding 0.1 ml of bromine to α,β -unsaturated compound (5) in methane dichloride drop wise in an ice bath under string for $1/2\text{ hr}$.^[29] The IR spectra for compound (5) showed absorption at 1695 cm^{-1} for ($\text{C}=\text{O}$) group and at 1125 cm^{-1} for ($\text{C}-\text{O}-\text{C}$) group. The ^1H NMR for dibromide showed a singlet signal at 8.56 ppm for C4 and two doublet at 7.92 and $7.48\text{--}7.49$ for C5, C7 and multiplet band at $4.20\text{--}4.32\text{ ppm}$ for furane and CHBr proton) table(1,2). Dimethyl sulfoxide in iodine is used as and iodinating agent for α,β -unsaturated ketone act as an oxidizing agent. The compound (6) is synthesized from adding one crystal of iodine to solution of compound (6) then acidified with two drops of H_2SO_4 ^[20]. Then the solution was refluxed for (1 hr).^[28], the IR spectra of compound (6) showed a strong absorption at 1685 cm^{-1} for ($\text{C}=\text{O}$) group and at 1565 cm^{-1} for ($\text{C}=\text{C}$) group and at 1120 cm^{-1} for ($\text{C}-\text{O}-\text{C}$) group. The ^1H NMR for compound (6) showed a doublet at 6.76 ppm for CH of ethylene and multiplet signal at $7.23\text{--}7.55\text{ ppm}$ for (benzene ring and CH of ethylene and C6 and CH furane) and singlet at 8.15 for C4 and two doublet at ($8.23, 8.52\text{ ppm}$) for ($\text{C}5,\text{C}7$) table(1,2).

Table 1. Physical and IR spectra data of compounds(I-VI).

Com. No.	Color	M.P.	Yield	IR data of compounds (cm ⁻¹)								
				C=O	CHO	C=N	C-Cl	Ar-H	OH	C-O-C	C-H	C=C
1	Pale Yellow	164	60%	1685	2720	1565	755	3035	--	--	--	--
2	Yellow	151	70%	1690	2720	1560	--	3065	3230	--	--	--
3	Yellow brawn	125-127	50%	1710	--	1555	--	3055	--	1120	2976	--
4	Pale Yellow	178-180	70%	1685	--	1550	--	3055	--	1120	--	1565
5	Brawn	203-205	70%	1695	--	1550	--	3060	--	1125	--	1580
6	Yellow	227-229	50%	1668	--	1560	--	3150	--	1125	--	--

Table 2. ¹H-NMR spectra data of compounds(I-VI).

Comp. No.	¹ H-NMR Chemical shift (DMSO-d ₆) δ , ppm
1	10.54(s,1H,CHO), 8.45(d,1H,C ₇), 7.86(s,1H,C ₄), 7.45(d,1H,C ₅), 7.27(t,1H,C ₆)
2	8.77(s,1H,CHO), 8.21(d,1H,C ₇), 8.05(s,1H,C ₄), 7.18(d,1H,C ₅), 7.02(t,1H,C ₆), 5.60(s,1H,OH)
3	2.99(s,3H,CH ₃), 6.99(s,1H,Furan), 7.15(t,1H,C ₇), 8.07(s,1H,C ₄), 8.21(m,1H,C ₅), 8.78(d,1H,C ₆)
4	6.53(d,CH, ethylene), 7.23-7.40(m,H-Furan,benzene ring), 7.30-7.32(d,1H,C ₇), 8.11(s,1H,C ₄), 8.67-8.68(m,1H,C ₅), 7.41-7.43(d,1H,C ₆)
5	7.20-7.32(m,2H,Furan,CH-Br), 7.56-7.92(m,6H, benzine ring & C ₆), 7.92(d,1H,C ₅), 8.56(s,1H,C ₄), 7.48-7.49(d,1H,C ₇)
6	7.23-7.55(m,8H, benzene ring, Furan, CH= & C ₆), 8.15(s,1H,C ₄), 8.23(d,1H,C ₅), 8.52(d,1H,C ₇)

Antibacterial activity

Anti-bacterial activity of new compounds was evaluated via paper disc-agar diffusion technique^[30] Ampicillin was used as standard for comparison. Anti-bacterial test was carried out by agar diffusion technique^[29,30] by measuring the diameter of zone of inhibition against test organisms at the end of (24h) at 37°C.

The compounds 3 and 6 showed inhibitory value against *Staphylococcus aureus* ranging at (18, 16) mm respectively and these value were less than the value of control factor antibiotics while compounds (4, 5) showed high inhibitory activity against the same bacteria the value of which was (20) mm which was very similar to an antibacterial effect of antibiotic ampicillin. The compounds (3, 4, 5) showed inhibitory value against the Gram negative bacteria *E. Coli* which ranged in value (16, 14) mm respectively , these values less than the control factor (18) mm while the compound (6) showed high inhibitory activity against *E. ccoli*. The compounds (3, 4 , 5 , 6) showed inhibitory value against *staphylococcus epidermis* (18, 18, 16, 12) respectively and these values were less than the value of control while all compound showed inhibitory vale against *proteus vagaries* less than the value of control. The results of biological tests are shown in **Table (3)**.

Table 3. Anti-bacterial activity data for compounds.

Compounds NO	Staphylococcus aurous	Staphylococcus epidermise	E.coli	Proteus vagaries
Ampicillin	20	20	18	18
Compound 3	18	18	16	12
Compound 4	20	18	16	14
Compound 5	20	16	14	12
Compound 6	16	12	18	14

4. Conclusion

The present study successfully synthesized a series of novel furor 2,3-b derivatives, including chalcone, dibromide, and iodo chalcone compounds, through multi-step reactions involving Vilsmeier–Haack condensation and Claisen–Schmidt condensation. The synthesized compounds were structurally confirmed by IR and ¹H NMR spectroscopy. Biological screening revealed that compounds (3–6) exhibited moderate antibacterial activity against both Gram-positive and Gram-negative bacteria, indicating their potential as promising scaffolds for future antimicrobial drug development.

Acknowledgment

Authors are grateful to the Researchers Supporting Project (ANUI2026SCI101), Alnoor University, Mosul, Iraq, the researchers also thank the University of Mosul and Northern Technical University for the support provided in the field of laboratory work.

Conflict of interest

The authors declare no conflict of interest.

References

1. Domínguez, J., Basante, W., Charris, J., & Riggione, F. (1996). Synthesis and activity of some quinolone derivatives against *Plasmodium falciparum* in vitro. *Farmaco* (Societa Chimica Italiana: 1989), 51(6), 407-412.
2. Matsumoto, J., Miyamoto, T., Minamida, A., Nishimura, Y., Egawa, H., & Nishimura, H. (1984). 1, 4-Dihydro-4-oxopyridinecarboxylic acids as antibacterial agents. 2. Synthesis and structure-activity relationships of 1, 6, 7-trisubstituted 1, 4-dihydro-4-oxo-1, 8-naphthyridine-3-carboxylic acids, including enoxacin, a new antibacterial agent. *Journal of medicinal chemistry*, 27(3), 292-301.
3. Banoon, S. R., & Ghasemian, A. (2022). The characters of graphene oxide nanoparticles and doxorubicin against HCT-116 colorectal cancer cells in vitro. *Journal of gastrointestinal cancer*, 53(2), 410-414.
4. Kuroda, T., Suzuki, F., Tamura, T., Ohmori, K., & Hosoe, H. (1992). A novel synthesis and potent antiinflammatory activity of 4-hydroxy-2 (1H)-oxo-1-phenyl-1, 8-naphthyridine-3-carboxamides. *Journal of medicinal chemistry*, 35(6), 1130-1136.
5. PL, F. (1998). Mori C. Badawneh M. Calderone V. Calzolari L. Loffredo T. Martinotti E. Saccomanni G. *Eur. J. Med. Chem.*, 33, 383.
6. Reddy, C. N., Prasad, P. R., & Sreedhar, N. Y. (2012). Electrochemical analysis of zincard in pharmaceutical and biological samples. *Indonesian Journal of Pharmacy*, 23(3), 147-155.
7. Bohm, B. A. (1998). *Introduction to flavonoids* (pp. xi+-503).
8. Prakash, O., Kumar, A., Sadana, A., Prakash, R., Singh, S. P., Claramunt, R. M., ... & Elguero, J. (2005). Study of the reaction of chalcone analogs of dehydroacetic acid and o-aminothiophenol: synthesis and structure of 1, 5-benzothiazepines and 1, 4-benzothiazines. *Tetrahedron*, 61(27), 6642-6651.
9. Prasad, Y. R., Rao, A. L., Prasanna, L., Murali, K., & Kumar, P. R. (2005). Synthesis and antidepressant activity of some 1, 3, 5-triphenyl-2-pyrazolines and 3-(2"-hydroxy naphthalen-1"-yl)-1, 5-diphenyl-2-pyrazolines. *Bioorganic & medicinal chemistry letters*, 15(22), 5030-5034.
10. Ramesh, S., Gaddam, V., & Nagarajan, R. (2010). A flexible approach to the chromenoquinolines under copper/Lewis acid catalysis. *Synlett*, 2010(05), 757-760.
11. Ningaiah, S., & Bhadregowda, D. G. (2024). Synthesis of chalcones derivatives and their biological potential: A review of nitrogen-containing heterocyclic scaffolds. *ACS Omega*, 9(14), 15891–15908. doi.org.
12. Dave, S. S., Ghatole, A. M., Rahatgaonkar, A. M., Chorghade, M. S., Chauhan, P. M. S., & Srivastava, K. (2009). Experimental and computational evaluation of new quinolinyl chalcones as potent antiplasmodial agents. *Indian journal of chemistry. Section B, Organic including medicinal*, 48(12), 1780.
13. Mistry, P. T., & Desai, K. R. (2021). Synthesis and antibacterial evaluation of some novel chalcones bearing 1,8-naphthyridine moiety under conventional and microwave irradiation. *Journal of Heterocyclic Chemistry*, 58(4), 932–941..
14. Rao, K. S., & Reddy, V. R. (2023). Development of a novel 1,8-naphthyridine–heterocycle hybrid derivatives: Synthesis, spectral characterization, and antimicrobial profiling. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 285, Article 121854.
15. Sadeek, G. T., Saeed, Z. F., & Saleh, M. Y. (2023). Synthesis and pharmacological profile of hydrazide compounds. *Research Journal of Pharmacy and Technology*, 16(2), 975-982.

16. Ruwaidah, H. A., & Dheefaf, F. S. (2023). Synthesis of novel heterocyclic compounds via cyclization reactions of Schiff bases. *Journal of the University of Baghdad for Science*, 20(3), 892–904.
17. Rani, P., & Pathak, D. P. (2024). Design, synthesis, and biological assessment of indole-carboxylic acid conjugates with promising cytotoxic and antimicrobial profiles. *Bioorganic Chemistry*, 142, Article 103984. doi.org
18. Ali, R. T., Mohammedthaji, N. H., & Al-Niemi, K. I. (2022). Study of Isothermal, Kinetic and Thermodynamic Parameters of Adsorption of Glycolic Acid by a Mixture of Adsorbent Substance with ab-Initio Calculations. *Egyptian Journal of Chemistry*, 65(6), 489-504.
19. Ghoneim, A. A., Elbargisy, R. M., & Manoer, A. (2020). Design and synthesis of heterocyclic Compounds from 1, 4-diacetylbenzene with Expected Antimicrobial Activity. *Egyptian Journal of Chemistry*, 63(8), 2901-2910.
20. Wang, L., Zhang, J., & Wang, Y. (2021). $\text{(Fe}_3\text{O}_4\text{@SiO}_2\text{-SO}_3\text{H)}$ nanocomposites: An efficient magnetically separable solid acid catalysts for an esterification reaction. *Micro & Nano Letters*, 12(4), 235–241. doi.org [1, 2]
21. Zolfigol, M. A., Khazaei, A., & Moosavi-Zare, A. R. (2021). Functionalized multi-walled carbon nanotubes as highly efficient heterogeneous solid acid catalysts in green organic transformations. *Journal of Nanoscience and Nanotechnology*, 21(4), 2134–2145.
22. Hasb, A. M., & Saeed, N. H. (2025). CHLORINATION OF N-BENZOYL VALINE BY SODIUM N-CHLORO-PARA-TOLUENE SULFONYL AMIDE (CAT) HYDROCHLORIC ACIDIC: A KINETIC AND MECHANISM STUDY. *Kimya Problemleri*, 23(2), 239-248.
23. Al-Adilee, K. J., & Al-Rubaie, L. A. (2020). Spectrophotometric determination of cefotaxime sodium via diazotization coupling reaction in pure and pharmaceutical samples. *Journal of Global Pharma Technology*, 12(6), 188–197.
24. Ali, A. H., Saleh, M. Y., & Owaid, K. A. (2023). Mild Synthesis, Characterization, and Application of some Polythioester Polymers Catalyzed by Cetrime Ionic Liquid as a Green and Eco-Friendly Phase-Transfer Catalyst. *Iranian Journal of Catalysis*, 13(1).
25. Banoon, S., Ali, Z., & Salih, T. (2020). Antibiotic resistance profile of local thermophilic *Bacillus licheniformis* isolated from Maysan province soil. *Comunicata Scientiae*, 11, e3291-e3291.
26. Saied, S. M., Mohammed, S. J., Khaleel, B. T., & Saleh, M. Y. (2021). Comparative studies between conventional techniques and green chemistry to synthesis of novel piperidinium salts ionic liquids (PBSILs).
27. Mohammed Abdulrazaq Mahdy, & Noor H. M. Saeed. (2025). A comparative study of the theoretical and practical values of ionization constants for amino acid derivatives in the neutral and anionic states. *Applied Chemical Engineering*, 8(4), ACE-5796.
28. Saeed, N. H., Ali, R. T., & Saied, S. M. (2025). Computational study on the Estramustine (EMCYT) and its active metabolites anticancer drugs 17- α -estradiol and nornitrogen. *Results in Chemistry*, 102747.
29. Rajesh kumar S. and Malavkodi, *Bioinorganic Chemistry and Applications*.(2014).
30. Carrod I.P. and Grady F.D. *Antibiotic and Chemotherapy*, rded. Churchill, livikgestons Edinburgh, 477(1972).